



## AccuStart™ II GelTrack™ PCR SuperMix (2X)

Cat No.	95136-100	Size:	100 x 25 µL reactions (1 x 1.25 mL)
	95136-500		500 x 25 µL reactions (5 x 1.25 mL)
	95136-04K		4000 x 25 µL reactions (1 x 50 mL)

Store at -25°C to -15°C

### Description

AccuStart II GelTrack PCR SuperMix is a 2X concentrated, ready-to-use reaction cocktail for routine PCR amplification of DNA fragments up to 4 kb followed by analysis on agarose gels. It contains all components, except primers and template. AccuStart II GelTrack PCR SuperMix simplifies reaction assembly, improves assay reproducibility, and reduces the risk of contamination. The supermix includes electrophoresis-tracking dyes that migrate at approximately 4 kb and 50 bp to allow direct loading of PCR product on agarose gels following amplification. AccuStart Taq DNA polymerase in the master mix is inactivated with monoclonal antibodies that bind the polymerase and keep it inactive prior to the initial PCR denaturation step. Upon heat activation (1 minute at 94°C), the antibodies denature irreversibly, releasing fully active, unmodified Taq DNA polymerase. This enables specific and efficient primer extension with the convenience of room temperature reaction assembly.

### Components

AccuStart II GelTrack PCR SuperMix (2X) 2X mix containing optimized concentrations of MgCl<sub>2</sub>, dNTPs, reaction buffer, AccuStart Taq DNA Polymerase, AccuStart Taq antibodies, stabilizers and gel loading dyes.

### Storage and Stability

Store components in a constant temperature freezer at -25°C to -15°C upon receipt.

Repeated freezing and thawing does not impair product performance.

For lot specific expiry date, refer to package label, Certificate of Analysis or Product Specification Form.

### Reaction Assembly

Component	Volume for 25-µL rxn.	Final Concentration
AccuStart II GelTrack PCR SuperMix (2X)	12.5 µL	1x
Forward primer	variable	100 – 500 nM
Reverse primer	variable	100 – 500 nM
Nuclease-free water	variable	
DNA Template	5 – 10 µL	Variable
Final Volume (µL)	25 µL	

**Note:** For smaller or larger reaction volumes (10 to 50 µL), scale all components proportionally.

### Reaction Protocol

Incubate the completed reaction mix in thermal cycler as follows:

Initial denaturation:	94°C, 1 to 3 min
PCR cycling (20 – 40 cycles:)	94°C, 10 to 30 s
	55 – 65°C, 15 to 30s
	68 – 72°C, 1 min per kb of product length
Hold	4°C until processed for analysis

Full activation of Taq DNA polymerase occurs within 30 seconds at 94°C. Complete denaturation of dsDNA target is important for efficient PCR amplification and may require different initial denaturation times depending on the properties of a given target sequence.

### Quality Control

Kit components are free of contaminating DNase and RNase. AccuStart II GelTrack PCR SuperMix is functionally tested for amplification of a 4-kb fragment from a single-copy gene in human genomic DNA.

### Limited Label Licenses

Use of this product signifies the agreement of any purchaser or user of the product to the following terms:

1. The product may be used solely in accordance with the protocols provided with the product and this manual and for use with components contained in the kit only. QIAGEN Beverly, Inc. grants no license under any of its intellectual property to use or incorporate the enclosed components of this kit with any components not included within this kit except as described in the protocols provided with the product, this manual, and additional protocols available at [www.quantabio.com](http://www.quantabio.com). Some of these additional protocols have been provided by QuantaBio product users. These protocols have not been thoroughly tested or optimized by QIAGEN Beverly, Inc.. QIAGEN Beverly, Inc. neither guarantees them nor warrants that they do not infringe the rights of third-parties.
2. Other than expressly stated licenses, QIAGEN Beverly, Inc. makes no warranty that this kit and/or its use(s) do not infringe the rights of third-parties.
3. This kit and its components are licensed for one-time use and may not be reused, refurbished, or resold.
4. QIAGEN Beverly, Inc. specifically disclaims any other licenses, expressed or implied other than those expressly stated.
5. The purchaser and user of the kit agree not to take or permit anyone else to take any steps that could lead to or facilitate any acts prohibited above. QIAGEN Beverly, Inc. may enforce the prohibitions of this Limited License Agreement in any Court, and shall recover all its investigative and Court costs, including attorney fees, in any action to enforce this Limited License Agreement or any of its intellectual property rights relating to the kit and/or its components.

©2018 QIAGEN Beverly Inc. 100 Cummings Center Suite 407J Beverly, MA 01915

QuantaBio brand products are manufactured by QIAGEN, Beverly Inc.

Intended for molecular biology applications. This product is not intended for the diagnosis, prevention or treatment of a disease.

AccuStart and GelTrack are trademarks of QIAGEN Beverly, Inc..