



repliQa HiFi ToughMix®

Cat No.	95200-025	Size:	25 x 50 µL reactions (1 x 0.625 mL)
	95200-100		100 x 50 µL reactions (1 x 2.50 mL)
	95200-500		500 x 50 µL reactions (1 x 12.50 mL)

Store at -25°C to -15°C

Description

The repliQa HiFi ToughMix is a unique, next generation 2x master mix that has 90x higher fidelity compared to *Taq* polymerase. The ToughMix has extreme speed, with extension times as fast as 1-10 sec/kb depending on target length. Additionally, the ToughMix has long range amplification properties as it can amplify fragments up to 24 kb from complex genomic DNA templates or fragments up to 40 kb from virus DNA templates such as *Escherichia virus Lambda* DNA.

The ToughMix is formulated with a genetically modified DNA polymerase coupled with hot start antibodies. It has 5' → 3' polymerase activity, 3' → 5' exonuclease activity, and generates blunt-ended products while providing the ability to amplify through uracils and primers containing inosine or uracil. It is *Tough Tested*, and is tolerant to multiple PCR inhibitors.

Components

repliQa HiFi ToughMix	2x reaction buffer containing optimized concentrations of MgCl ₂ , dNTPs and proprietary formulated HiFi polymerase, hot start antibodies and ToughMix chemistry
-----------------------	---

Storage and Stability

Store kit components in a constant temperature freezer at -25°C to -15°C protected from light upon receipt. For lot specific expiry date, refer to package label, Certificate of Analysis or Product Specification Form.

Guidelines for PCR

- The design of highly specific primers is a critical parameter for successful PCR. The use of computer aided primer design programs is encouraged in order to minimize the potential for internal secondary structure. For best results, primer size should be limited to 22 - 35 bp with a melting point of at least 63°C. Ideal GC-content of the primers is 45-60%. A final concentration of 300 nM each primer is effective for most applications. Primers with inosine (dI) and uracil (dU) are acceptable.
- Preparation of a reaction cocktail is recommended to reduce pipetting errors and maximize assay precision. Assemble the reaction cocktail with all required components except sample template and dispense equal aliquots into each reaction tube. Add the DNA template to each reaction as the final step. Addition of samples as 2 to 5-µL volumes will improve assay precision.
- Suggested input quantities of template are: genomic DNA ≤ 200 ng; plasmid DNA ≤ 50 ng; cDNA ≤ 750 ng.
- After sealing each reaction, vortex gently to mix contents. Centrifuge briefly to collect components at the bottom of the reaction tube.
- Longer targets may require a higher primer melting temperature of at least 65°C, and a lower primer concentration of 150 nM.

Reaction Assembly

Component	Volume for 50 µL rxn	Final Concentration
repliQa HiFi ToughMix (2X)	25 µL	1x
Forward primer	variable	300 nM
Reverse primer	variable	300 nM
Nuclease-free water	variable	
Template	2 – 5 µL	variable
Final Volume (µL)	50 µL	

Note: For smaller or larger reaction volumes scale all components proportionally.



PCR Cycling Protocol

Initial denaturation and final extension steps are typically not required with repliQa HiFi ToughMix.

The 10 s at 98°C during cycling are sufficient to fully activate the HotStart mechanism. For longer fragments, (>10 kb), an initial denaturation of 98°C for 30 s can be added to facilitate denaturation of the DNA template.

PCR cycling (25 - 45 cycles):

*We recommend using 2-step cycling first. If you are having difficulty, we suggest trying 3-step cycling.

2-Step Cycling*	3-Step Cycling
98°C, 10 s	98°C, 10 s
	(T _m -5) °C, 5 s
68°C	68°C
≤ 1 kb: 1 sec	≤ 1 kb: 1 sec
1 ~ 10 kb: 5 sec/ kb	1 ~ 10 kb: 5 sec/ kb
≥ 10 kb: 10 sec/ kb	≥ 10 kb: 10 sec/ kb

Quality Control

Kit components are free of contaminating DNase and RNase. 2x repliQa HiFi ToughMix is functionally tested for amplification of a 4-kb fragment from a single-copy gene in a human genomic DNA.

Nuclease Assay:

DNase: DNase activity must be below the detectable limits of 100 pg DNase I equivalent as assayed using a fluorogenic substrate following a 1 hour incubation at 37°C with each kit component at 1x concentration.

RNase: RNase activity must be below the detectable limits of 1 pg RNase A equivalent as assayed using a fluorogenic substrate following a 1 hour incubation at 37°C with each kit component at 1x concentration.

4.1 kb PCR Functional Assay: Negative control must be free of visible product with a single band at ~4.1 kb visible from 35 cycles of PCR using 20 ng human genomic DNA.

Limited Label Licenses

Use of this product signifies the agreement of any purchaser or user of the product to the following terms:

1. The product may be used solely in accordance with the protocols provided with the product and this manual and for use with components contained in the kit only. QIAGEN Beverly, Inc. grants no license under any of its intellectual property to use or incorporate the enclosed components of this kit with any components not included within this kit except as described in the protocols provided with the product, this manual, and additional protocols available at www.quantabio.com. Some of these additional protocols have been provided by Quantabio product users. These protocols have not been thoroughly tested or optimized by QIAGEN Beverly, Inc.. QIAGEN Beverly, Inc. neither guarantees them nor warrants that they do not infringe the rights of third-parties.
2. Other than expressly stated licenses, QIAGEN Beverly, Inc. makes no warranty that this kit and/or its use(s) do not infringe the rights of third-parties.
3. This kit and its components are licensed for one-time use and may not be reused, refurbished, or resold.
4. QIAGEN Beverly, Inc. specifically disclaims any other licenses, expressed or implied other than those expressly stated.
5. The purchaser and user of the kit agree not to take or permit anyone else to take any steps that could lead to or facilitate any acts prohibited above. QIAGEN Beverly, Inc. may enforce the prohibitions of this Limited License Agreement in any Court, and shall recover all its investigative and Court costs, including attorney fees, in any action to enforce this Limited License Agreement or any of its intellectual property rights relating to the kit and/or its components.

©2020 QIAGEN Beverly Inc. 100 Cummings Center Suite 407J Beverly, MA 01915

Quantabio brand products are manufactured by QIAGEN, Beverly Inc.

Intended for molecular biology applications. This product is not intended for the diagnosis, prevention or treatment of a disease.

Trademarks

ToughMix is a registered trademark of QIAGEN Beverly, Inc.