

Product Information	
<b>qScript® cDNA SuperMix</b>	
Part Number	95048-500
Number of Reactions	500 Reactions
Reaction Size	20 µL
Storage Temperature	-25°C to -15°C
Lot Number	026419
Reference Number	040218
Expiration Date	08/31/2019

**Product Description:**

qScript cDNA SuperMix is a 5X concentrated, sensitive, and easy-to-use 1-tube reagent for first-strand cDNA synthesis that combines a highly-modified RNase H<sup>+</sup> mutant of M-MLV together with ribonuclease inhibitor protein (RIP) in a rigorously optimized formulation for real-time qPCR applications. The stabilized SuperMix formulation has been rigorously optimized to deliver sensitive, linear assay performance across a spectrum of relative abundance and input RNA (10pg - 1ug). qScript cDNA SuperMix reagent performance is unaffected by repetitive freeze/thaw cycling (>20X), conferring greater ease-of-use and consistent assay performance. Oligo (dT) and random primers are pre-blended in a precise ratio to provide equal representation of 5' and 3'-sequences for accurate gene expression quantification.

**Component Part Numbers:**

84035 qScript cDNA SuperMix, 1000 µL

Product Specifications			
95048			
Assay	cDNA SuperMix Functional qPCR Assay	DNase	RNase
Result	Pass	Pass	Pass

**Quality Control Analysis and Specifications:**
**Nuclease Assay:**

**DNase:** DNase activity must be below the detectable limits of 100 pg DNase I equivalent as assayed using a fluorogenic substrate following a 1 hour incubation at 37°C with each kit component at 1X concentration.

**RNase:** RNase activity must be below the detectable limits of 1 pg RNase A equivalent as assayed using a fluorogenic substrate following a 1 hour incubation at 37°C with each kit component at 1X concentration.

**cDNA SuperMix Functional qPCR Assay:** First-strand synthesis is performed on a 10-fold serial dilution over 6 orders of dynamic range (1 µg to 1 pg) using a Universal Reference total RNA preparation. One tenth of each first strand reaction is used as template for real-time PCR of a reference gene in duplicate reactions. Cq standard curve analysis must have coefficient of determination ( $r^2$ )  $\geq 0.990$  with a slope between  $-3.20$  and  $-3.70$ .

**Limitations of Use**

QuantaBio and Ultraplex are registered trademarks of QIAGEN Beverly, Inc. Quanta Biosciences, qScript, Geltrack, ToughMix, PerfeCta, and Fastmix are registered trademarks of Quanta BioSciences Inc. Extracta, AccuStart, AccuMelt, and Accuvue are trademarks of Quanta BioSciences Inc. Applied Biosystems, StepOne, StepOnePlus and ROX are trademarks of Thermo Fisher Scientific and or its subsidiaries. Please contact QIAGEN-Beverly for more information.

This product was developed, manufactured, and sold for *in vitro* use only. The product is not suitable for administration to humans or animals. SDS sheets relevant to this product are available upon request.

100 Cummings Center, Suite 407J, Beverly, MA 01915 • Ph (888) 927-7027 • Fax (978) 867-5724 • [www.QuantaBio.com](http://www.QuantaBio.com) FMWI016.2 Rev 01