

Product Information	
qScript® cDNA SuperMix	
Part Number	95048-100
Number of Reactions	100 Reactions
Reaction Size	20 µL
Storage Temperature	-25°C to -15°C
Lot Number	66259580
Reference Number	022123
Expiration Date	02/28/2026

**Product Description:**

qScript cDNA SuperMix is a 5X concentrated, sensitive, and easy-to-use 1-tube reagent for first-strand cDNA synthesis that combines a highly-modified RNase H<sup>+</sup> mutant of M-MLV together with ribonuclease inhibitor protein (RIP) in a rigorously optimized formulation for real-time qPCR applications. The stabilized SuperMix formulation has been rigorously optimized to deliver sensitive, linear assay performance across a spectrum of relative abundance and input RNA (10pg - 1ug). qScript cDNA SuperMix reagent performance is unaffected by repetitive freeze/thaw cycling (>20X), conferring greater ease-of-use and consistent assay performance. Oligo (dT) and random primers are pre-blended in a precise ratio to provide equal representation of 5' and 3'-sequences for accurate gene expression quantification. For gene-specific priming (GSP) or two-step RT-PCR of RNA exceeding 1kb total length, see our qScript Flex cDNA Kit.

**Component Part Numbers:**

84034 qScript cDNA SuperMix, 400µL

Product Specifications			
95048			
Assay	cDNA SuperMix Functional qPCR Assay	DNase	RNase
Result	Pass	Pass	Pass

**Quality Control Analysis and Specifications:**
**Nuclease Assay:**

**DNase:** DNase activity must be below the detectable limits of 100 pg DNase I equivalent as assayed using a fluorogenic substrate following a 1 hour incubation at 37°C with each kit component at 1X concentration.

**RNase:** RNase activity must be below the detectable limits of 1 pg RNase A equivalent as assayed using a fluorogenic substrate following a 1 hour incubation at 37°C with each kit component at 1X concentration.

**cDNA SuperMix Functional qPCR Assay:** First-strand synthesis is performed on a 10-fold serial dilution over 6 orders of dynamic range (1 µg to 1 pg) using a Universal Reference total RNA preparation. One tenth of each first strand reaction is used as template for real-time PCR of a reference gene in duplicate reactions. Cq standard curve analysis must have coefficient of determination ( $r^2$ )  $\geq 0.990$  with a slope between -3.20 and -3.70.

**Limitations of Use**

QuantaBio, UltraPlex, qScript, GelTrack, ToughMix, PerfeCta, and FastMix are registered trademarks of QuantaBio, LLC. Applied Biosystems, StepOne, StepOnePlus and ROX are trademarks of Thermo Fisher Scientific and or its subsidiaries. Please contact QuantaBio for more information.

This product was developed, manufactured, and sold for *in vitro* use only. The product is not suitable for administration to humans or animals. SDS sheets relevant to this product are available upon request.

100 Cummings Center, Suite 407J, Beverly, MA 01915 • Phone +1 888-959-5165 • [www.quantabio.com](http://www.quantabio.com) FMWI-016.2 Rev.03